

# **Climate change and its impacts in the ASEAN region**

**Second China-ASEAN Academy on Ocean  
Law and Governance**

**6-17 Nov 2016, Haikou, Hainan**

**Poh Poh Wong**

**Email: [wong3921@gmail.com](mailto:wong3921@gmail.com)**

**Website: [coast4all](http://coast4all.com)**

# Outline

- 1. IPCC**
- 2. ASEAN**
- 3. Climate change**
- 4. Coast**
- 5. Freshwater**
- 6. Food security**
- 7. Human settlements**
- 8. Adaptation**

# 1. IPCC

# IPCC

- Created in **1988** by WMO and UNEP. Has **195 governments** that commission assessments performed by the international community on the state of human knowledge of climate & climate change.
- Role : to assess on a comprehensive, objective, open & transparent basis the scientific, technical & socio-economic information relevant to **understanding scientific basis of risk of human-induced climate change, its potential impacts & options for adaptation & mitigation.**
- IPCC assessments : scientific basis for governments at all levels to develop climate related policies, & they **underlie negotiations at** the UN Climate Conference (**UNFCCC**, United Nations Framework Convention on Climate Change).



# WGs and Task Force

- **WGI** : assesses physical **scientific aspects** of climate system & climate change.
- **WGII** : assesses **vulnerability** of socio-economic & natural systems to climate change & **options for adapting** to it.
- **WGIII** : assesses **options for mitigation** climate change through limiting or preventing GHG emissions & enhancing activities to remove them.
- **Task Force** : to develop & refine internationally-agreed methodology & software for calculation & reporting **GHG emissions**.


# WGII AR5

- **242 Lead Authors (about 60 are CLAs) & 66 Review Editors from 70 countries** : selection criteria include expertise, differing viewpoints & perspectives, geographic balance, gender balance, & ensuring involvement of new experts in accordance to agreed-upon IPCC guidelines. Supplemented by **436 Contributing Authors from 54 countries**.
- Undergone two **extensive reviews** : totaling over 50,000 comments from **1,729 expert reviewers from 84 countries & 49 governments**.
- **Large knowledge base**. Substantial larger knowledge base : **30 chapters (from 20 in AR4); additional 4 on adaptation; 4 on livelihoods & poverty, human security, urban & rural areas; 2 on oceans. Two parts; part 1 on global & sectoral aspects; part 2 on regions. Over 12,000 scientific references.**

# AR5 reports



# AR5 RCPs



- **Representative concentration pathways (RCPs)** are scenarios that include time series of emissions & concentrations of the full suite of GHGs and aerosols and chemically active gases, as well as land use/land cover. *Representative* signifies that each RCP provides only one of many possible scenarios that would lead to the specific radiative forcing characteristics.
- **4 RCPs** used in WGII : RCP8.5 (high), RCP6.0 and RCP4.5 (intermediate), RCP2.6 (low) – named after level of radiative forcing reached or stabilized in 2100.




# Projected MSLR

Emission scenario	Representative Concentration Pathway (RCP)	2100 CO <sub>2</sub> concentration (ppm)	Mean sea level rise (m)	
			2046–2065	2100
Low	2.6	421	0.24 [0.17–0.32]	0.44 [0.28–0.61]
Medium low	4.5	538	0.26 [0.19–0.33]	0.53 [0.36–0.71]
Medium high	6.0	670	0.25 [0.18–0.32]	0.55 [0.38–0.73]
High	8.5	936	0.29 [0.22–0.38]	0.74 [0.52–0.98]


Emission scenario	Mean sea level rise (m)		
	2200	2300	2500
Low	0.35–0.72	0.41–0.85	0.50–1.02
Medium	0.26–1.09	0.27–1.51	0.18–2.32
High	0.58–2.03	0.92–3.59	1.51–6.63

# Managing risks

- **Risk** : potential for consequences where something of human value (including humans themselves) is at stake and where the outcome is uncertain.  
**Risk = hazard x vulnerability x exposure.**
- **Risk management** : Plans, actions, or policies implemented to reduce the likelihood &/or consequences of risks or to respond to consequences.



# Characterizing risks



# 2. ASEAN

# ASEAN and SE Asia




## Southeast Asia (12)

- Brunei
- Indonesia
- Lao People's Democratic Republic
- Malaysia
- Myanmar
- Papua New Guinea
- The Philippines
- People's Republic of Cambodia
- Singapore
- Thailand
- Timor-Leste
- Vietnam

# Vulnerability

- **One of most vulnerable regions** (others being South Asia & small island states). Due to long coastlines, high concentration of population & economic activity in coastal areas, & heavy reliance on agriculture, natural resources, & forestry.
- **Already affected by climate change** : increasing frequency & intensity of extreme weather events such as heat waves, droughts, floods & tropical cyclones in recent decades. Exacerbating water shortages, constraining agricultural production & threatening food security, causing forest fires & coastal degradation, & increasing health risks.
- **Some highlights** : impacts & adaptation options in four key areas - coasts, freshwater, food security, human settlements.





(EEPSEA 2009)

# **3. Climate change**



# Regional features

- **Complex range** of terrains & land–sea contrasts.
- With long coastlines therefore include some **influence of the ocean**.
- Several **large-scale phenomena** influence the climate of this region : monsoons, IOD, ENSO, TC, MJO.
- Climate variability & trends **differ vastly** across the region & between seasons. **Strong seasonality** in change is observed.

# Monsoons

- **East Asian monsoon** : monsoonal flow that carries moist air from Indian Ocean & Pacific Ocean to East Asia, affecting most of SE Asia.
- **Projected monsoon-related interannual rainfall variability** will increase in future. Future increase in **precipitation extremes** related to monsoon is very likely.

# Indian Ocean Dipode (IOD)

- **IOD (Indian Niño)** : an irregular oscillation of sea-surface temperatures in which western Indian Ocean becomes alternately warmer & then colder than eastern part of the ocean.
- **Projected zonal (east–west) pattern of change** : with reduced warming & decreased precipitation in the east, & increased warming & increased precipitation in the west, directly influencing SE Asian precipitation.
- Impact **more prominent in eastern Indonesia.**



# El Niño Southern Oscillation (ENSO)

- **Warming of ocean surface** or above-average sea surface temperatures in either central & eastern tropical Pacific Ocean. ENSO influence is predominant in East Malaysia & areas east of it. Reducing rainfall over Indonesia.
- Reduction in mean precipitation & enhanced warming if El Niño events become more frequent &/or intense.
- **Projected dominant mode of interannual variability** in the future; however, any specific projected change in its variability in the 21st century remains low.

# Tropical cyclones

- **Rapidly rotating storm system.** Increasing frequency of extreme events reported in northern parts of SE Asia, decreasing trends reported in Myanmar.
- Projected low confidence in region-specific projections of frequency & intensity.  
**Projected increase** in extreme precipitation near centres of tropical cyclones.

# Tropical cyclones (typhoons)



(Rohde)

# Examples

- **Tropical cyclones** (typhoons) : (frequency & intensity of cyclones not fully known); storm surge sensitive to tropical cyclones; precipitation be more extensive where tropical cyclones make landfall.
- **Monsoons** : increases in precipitation extremes.

## Worst Storms in Southeast Asia



# Madden-Julian Oscillation (MJO)

- Largest element of the **intraseasonal (30- to 90-day) variability in the tropical atmosphere**. Unlike a standing pattern like the ENSO, MJO is a traveling pattern that propagates eastward. Associated with enhanced rainfall in Indonesia during northern winter & SE Asia during northern summer.
- Due to poor ability of models to simulate it & its sensitivity to ocean warming patterns, future **projections of regional climate extremes in SE Asia** are therefore **highly uncertain** when associated with the MJO.




# Observed climate change

- **Temperature : increasing** at a rate of 0.14°C to 0.20°C per decade since 1960s, coupled with rising number of hot days & warm nights, & decline in cooler weather.
- **Rainfall** : annual total wet-day rainfall **increased** by 22 mm per decade, while rainfall from extreme rain days increased by 10 mm per decade, but climate variability & trends differ vastly across the **region** & **between seasons**.
- While an **increasing** frequency of **extreme events** has been reported in the northern parts of SE Asia, **decreasing** trends in such events are reported in Myanmar.


# Temperature change

- The **median increase in temperature over land** ranges from  $0.8^{\circ}\text{C}$  in RCP2.6 to  $3.2^{\circ}\text{C}$  in RCP8.5 by the end of this century (2081–2100).

Temperature change Southeast Asia (land) December-February




Temperature change Southeast Asia (sea) December-February



# Temp. mean changes for 2081-2100


- **Warming** is very likely to continue with substantial sub-regional variations.




# Precipitation change

- Moderate increase in precipitation is projected : 1% in RCP2.6 increasing to 8% in RCP8.5 by 2100.**


Precipitation change Southeast Asia (land) April-September




Precipitation change Southeast Asia (land) October-March



Precipitation change Southeast Asia (sea) April-September






Precipitation change Southeast Asia (sea) October-March












# Precip. mean changes for 2081-2100

- Moderate increase in rainfall, except on Indonesian islands neighbouring southeast Indian Ocean.
- On **islands neighbouring the southeast** tropical Indian Ocean, **rainfall is projected to decrease** during July to November (the IOD prevalent season), consistent with a slower oceanic warming in the east than in the west tropical Indian Ocean despite little change in IOD.
- **Strong regional variations** expected because of terrain.



# Summary - climate change

Region/ region code	Trends in daytime temperature extremes (frequency of hot and cool days)		Trends in heavy precipitation (rain, snow)		Trends in dryness and drought	
	Observed	Projected	Observed	Projected	Observed	Projected
<b>Southeast Asia</b> <b>SEA, 24</b>	 Increases in hot days (decreases in cool days) for northern areas <sup>a</sup>   Insufficient evidence for Malay Archipelago <sup>a</sup>	 <i>Likely</i> increase in hot days (decrease in cool days) <sup>b</sup>	 Spatially varying trends, partial lack of evidence <sup>a</sup>	 Increases in most metrics over most (especially non- continental) regions. One metric shows inconsistent signals of change. <sup>b</sup>	 Spatially varying trends <sup>a</sup>	 Inconsistent signal of change <sup>b</sup>

(TS 2014)

# Question 1



- **What is the extent of knowledge of climate change in your country?**

# 4. Coasts




# Coastal systems

- Low-lying coasts support **mangrove forests**, most of world **peat swamp forests**.
- 40% of world's **coral reefs** in Asia, mostly in SE Asia.
- World's most diverse reef communities in the "**coral triangle**".
- Also widespread **seagrass beds**.



# Coastal risks



**Figure 5-1** | Climate, just as anthropogenic or natural variability, affects both climate and human related drivers. Risk on coastal systems is the outcome of integrating drivers' associated hazards, exposure, and vulnerability. Adaptation options can be implemented either to modify the hazards or exposure and vulnerability, or both.

# Projected impacts on coasts

- **Increasing SST & ocean acidification** : declines in coral-dominated reefs & other calcified marine organisms. On rocky shores warming & acidification are expected to lead to range shifts & changes in biodiversity.
- **SLR** : increase coastal flooding, coastal erosion & saltwater intrusion; mangroves, salt marshes & seagrass beds decline unless they receive sufficient fresh sediments to keep pace of SLR or retreat inland.
- **Cyclone intensification with SLR** : increase coastal flooding. In 2013, 176 million people in (Asian) region were affected by natural disasters such as flooding & cyclones, with 3.5 million displaced due to climate-related catastrophes.

# Acidification and coral reefs

- Continuation of current trends in sea surface **temperatures & ocean acidification** would result in large declines in coral-dominated reefs by mid-century.
- **Coral reefs** will be most vulnerable marine ecosystem with little scope for adaptation. Warming & acidification will lead to coral bleaching, mortality & decreased constructional ability. [Acidification highlighted in AR5 than in AR4, due to more information & two ocean chapters in AR5]
- Acidification also expected to have negative impacts on **other calcified marine organisms** with large & uncertain regional & local variations.

# SLR

- **Future rates of SLR** are expected to exceed those of recent decades, increasing coastal flooding, erosion, & saltwater intrusion into surface & groundwater.
- In the absence of other impacts, **coral reefs** may grow fast enough to keep up with rising sea levels; but **beaches** may erode & **mangroves, salt marshes, & seagrass** beds will decline, unless they receive sufficient fresh sediment to keep pace or they can move inland
- Coastal freshwater wetlands may be vulnerable to saltwater intrusion with rising sea levels, but in most river **deltas** local subsidence for nonclimatic reasons will be more important.
- Current trends in **cyclone frequency & intensity** are unclear, but combination of cyclone intensification & SLR could increase coastal flooding.

# RSLR

- **RSLR** at local scale can be **much larger** than projected GMSLR.
- **Non-climate change** local processes include subsidence, glacial isostatic adjustment, sediment transport, coastal development.
- Changes in storms & associated storm surges may further contribute to **sea level extremes**.
- Under present levels of global warming, already **committed to higher future SLR** above current levels.


# Non-climate factors

- Increase significantly in future due to **population growth, economic development, & urbanization.**
- Humans have been primary drivers of change in **coastal aquifers, lagoons, estuaries, deltas & wetlands.**
- Further exacerbation on coastal ecosystems from **excess input, changes in runoff & reduced sediment delivery.**



# Coastal zones at risk

## VISUALISING A WARMING WORLD COASTAL ZONES AND PRODUCTIVITY AT RISK IN SOUTHEAST ASIA



Designed by Laura Connor

Sources:  
Turn Down the Heat: Climate Extremes, Regional Impacts and the Case for Resilience, World Bank, 2013  
Reefs at Risk - Revisited, World Resources Institute, 2011  
Joint Typhoon Warning Center, 2011






# SLR vulnerability



(Jones et al 2012)

# SLR – 1 m SLR



# 5. Freshwater

# Freshwater resources

- **Importance** : massive population & heavy dependence of agricultural sector on precipitation, river runoff, & groundwater.
- **Non-climate drivers** : soaring populations, increasing per-capita domestic use, due to urbanization & thriving economic growth, & increasing use of irrigation.
- **Water scarcity** is expected to be a big challenge in many regions because of increasing water demand from population growth & consumption per capita with higher standards of living.

# Aquifers

- **Groundwater sources**, which are affordable means of high-quality water supply in cities of developing countries, are threatened due to over-withdrawals.
- **Aquifer levels have fallen** by 20 to 50 m in cities such as Bangkok & Manila. Drop in groundwater levels often results in land subsidence, which can enhance hazard exposure due to coastal inundation & SLR especially in settlements near the coast, & deterioration of groundwater quality.

# Projected impacts

- **Future water availability** differs substantially among river basins & seasons.
- **Low confidence in projections** of specifically how climate change will impact future precipitation on sub-regional scale, & in projections of how climate change might impact availability of water resources.
- Future projections suggest a decrease in river runoff in January in **Chao Phraya River** basin, Thailand.

# 6. Food security




# Key drivers

- Food production & food security are most **vulnerable to rising air temperatures.** Warmer temperatures could depress yields of major crops such as rice.
- **SLR will be a key issue** for many coastal areas as rich agricultural lands may be submerged & taken out of production.

# Agriculture - temperature

- **Current temperatures** : already approaching critical levels during susceptible stages of the rice plant in Myanmar/Thailand/Laos/Cambodia (March-June), Vietnam (April/August), Philippines (April/June), Indonesia (August).
- With **rising temperatures**, the process of rice development accelerates & reduces the duration for growth.


(USDA)




# Agriculture - SLR

- SLR : **key issue** for many coastal areas as rich agricultural lands may be submerged.
- **Threatens coastal & deltaic rice production** areas particularly in Mekong River Delta. E.g. about 7% of Vietnam's agriculture land may be submerged, decrease in Myanmar's rice yield due salt water intrusion.

# Mekong Delta – 0.3 m SLR on rice




SLR by 30 cm:  
50,000 ha affected ( of 1.8m ha)  
• 120,000 tonnes less rice (23m tonnes)



SLR 30 cm + dams + drought  
500,000 ha affected  
• 1,000,000 tonnes less rice


# Fisheries

- Concern centered on **rising water temperatures** & potential impacts of climate change on flow regimes, which affect the reproduction of many fish species.
- **Decline in marine productivity** in part due to vulnerability of coral reefs to both warming & ocean acidification.
- Climate change may lead to massive **redistribution of fisheries** catch potential with large declines in the tropics, particularly Indonesia.
- **Destructive** fisheries : human factor.



# Deltas – capture fisheries & aquaculture

- **Lower Mekong River basin** supports largest freshwater capture fishery in the world & would be negatively impacted.
- **Sea level rise** is expected to impact both capture fisheries & aquaculture production in river deltas : increasing salinity.
- Based on study of Yellow River delta, aquaculture in Mekong Delta may contribute to SLR – land subsidence caused by groundwater pumping (Higgins et al 2013).







# **7. Human settlements**



# Population centres in deltaic areas

- **Population centres in deltaic areas** : exposed to high degree of cumulative climate-related risk (all environmental & socioeconomic factors).



# Cities vulnerable to coastal flooding

- **By 2070 : SE Asian cities with most risk to coastal flooding include Ho Chi Minh City, Bangkok, Rangoon, Jakarta & Manila.**



Key South East Asian Agglomerations	Population (2005, in millions)	Projected Exposed Population (2070, in millions)	Local Sea Level Rise Projections in a 4°C World in 2070 (above 1986–2005)
Jakarta	13.2	2.2	66cm
Yangon	4.1	4.9	63cm
Manila	10.6	0.5	66cm
Bangkok	6.5	5.1	65cm
Ho Chi Minh City	5.0	9.2	65cm

Source: Population data from Hanson et al. (2011); SLR RCP8.5 (in this report).

(ADB 2013)

# Subsidence

- Groundwater withdrawal & surface drainage for urban development: resulting in **land subsidence** in Bangkok, Jakarta, Manila, Ho Chi Minh, Semarang, etc, increasing hazard exposure to coastal inundation & SLR.



# Question

- **Which sector of your country is most affected by climate change?**
- **And what adaptation measures are being planned?**

# 8. Adaptation

# Adaptation - freshwater

- Develop **adaptive/integrated water resource management** of the trade-offs balancing water availability against increasing demand.
- **Diversify water supply sources** including reuse & recycling, use of household-level water resources, e.g. roof water harvesting.
- **Reservoirs** partly mitigate seasonal differences & increase water availability for irrigation.
- **In coastal areas, desalination** of seawater or brackish water to supplement.
- **Integrated water management within river basins** to benefit countries.



# Adaptation – food security

- **Crop breeding** : for high temperature condition. Promising option. E.g. 9 salt-tolerant rice varieties in 2013 in Philippines.
- **Aquaculture : better management practices** for shrimp, pangasius catfish, brackish-water & nearshore aquaculture, marine cage culture (Thailand, Philippines, Indonesia, Vietnam).



Source - IRRI

(IRRI)



(BFAR)



# Adaptation – human settlements

- Focus on solely **adapting through physical infrastructure** in urban areas requires complementary adapting planning, management, governance & institutional arrangements to deal with uncertainty & unprecedented challenges.
- **Integrating DRR & CCA** in urban development, urban planning.

Urban development	<ul style="list-style-type: none"><li>• Integrate flood risk management into spatial planning to protect groundwater recharge zones and floodplains</li><li>• Adopt building codes and infrastructure standards to consider changes such as new flood return periods</li><li>• Prepare disaster preparedness plans that consider disruptions to services</li><li>• Develop measures that address the needs of vulnerable populations</li></ul>
Transport	<ul style="list-style-type: none"><li>• Redesign or relocate road facilities</li><li>• Add or redesign protective measures for road corridors and coasts</li><li>• Increase drainage for road facilities against projected increases in precipitation and erosion</li><li>• Ensure road access to hospitals and evacuation centers and distribution of relief where road infrastructure may be damaged during extreme events</li></ul>

(ADB 2013)

# Adaptation - coasts

- **Creation of MPAs** targetting areas where SST are projected to change least : increase resilience.
- **Hard coastal defenses**, such as sea walls, protect settlements at the cost of preventing adjustments by mangroves, salt marshes, & seagrass beds to rising sea levels.
- **Landward buffer zones** : for inland migration of mangroves & seagrasses.

# Adaptation and mitigation

- **Agroforestry practices** : carbon storage, decrease soil erosion, increase resilience against floods, landslides & drought, biodiversity benefits, reduce financial impact of crop failure, etc.
- Very high **carbon sequestration potential** of organic-rich soils in mangroves & peat swamp forests.
- **Sustainable cities** with fewer fossil-fuel driven vehicles (mitigation) & more trees & greenery.

# EBA - Ecosystem-based management


- **Emerging approach** using biodiversity & ecosystem services (supporting, provisioning, regulating, cultural) as part of overall strategy to adapt to adverse effects of climate change.
- **Part of broader portfolio** of adaptation measures – can be applied at different geographical scales & within various time frames.
- Is **cost-effective** to protect communities from climate change & extreme weather events.

# EBA for coasts and islands

- “Natural ecosystems such as coastal forests, coral reefs, **mangrove belts**, beach ridges, sand dunes or forested slopes are effective barriers against many types of natural disasters. Such reinforcements can be a cost-effective insurance against storm surges, tsunami and sea-level rise for coastal communities that cannot afford expensive infrastructural protection.” (*Planet Prepare* 2008: 63)

# Mangroves

- Have **special root systems** & may adapt to changes in sea level by growing upward in place, or by expanding landward or seaward.



Peg roots of  
*Avicennia*




Knee Roots of  
*Bruguiera*



Peg roots of  
*Sonneratia*



Prop roots of  
*Rhizophora*



© PP Wong



© PP Wong



© PP Wong

# Buffer to waves

- Publications on experiments & field studies done on effectiveness of mangroves (Mazda et al 2007).
- **Review by Cochard et al (2008)** summed up & put controversy to rest to some extent. Considerable buffer to waves up to 4 m high.
- **Reduce wave ht & energy** by 13% to 66% & surges by 50 cm for every km, as they pass through trees & exposed roots.

Ecosystem type	Dominant ecosystem processes	Dominant buffer composition	Approximate wave buffer effectiveness range				Expected tsunami energy exposure
			Normal waves	Storm waves	<4 m high tsunami	>8 m high tsunami	
(c) Mangrove forests	Biotic/physical	Biotic	▼~▼ <sup>1</sup>	▼~▼ <sup>1,2</sup>	▼~▼ <sup>2</sup>	▲~▼ <sup>2</sup>	■—■

Legend:


▲	Hazard amplification	▼ ▲	Slight effect (not evident, but measurable)	■	Small
▼	Hazard mitigation	▼	Moderate effect (evident, ~20-50% energy reduction)	■	Medium
X	No effect	▼	Considerable effect (~50-100% energy reduction)	■	High

(Cochard et al 2008)



# Keeps up with SLR

- Keeps up with SLR : **1 mm/annum.**
- “Intact and healthy mangrove systems can adapt to sea level rise; their growth **can accommodate to increases of 3.8 up to 9 mm per year** depending on local circumstances.”  
(Wetlands International)





(Alongi 2008)

# Large-scale modular planting (1)

- Large-scale planting using modular system to meet requirements of various coastal locations. **Modular system of planting & deployment** is comparable to LEGO® set on large scale.
- Ideally space-fitting shapes containing sediments with mangroves grown to various heights. Modules made of local materials or mixture of compressed sediments that become self-destructive & formed part of substrates. Nutrients & sediments added to growing mangroves in field (Wong 2011).
- Suitable **for variety of coasts** & not confined to muddy coasts.

# Large-scale modular planting (2)





# *Avicennia marina*

- **Widest latitudinal range**, ability to adapt to wide range of physical conditions, only mangrove to survive in arid areas.
- Present on both **seaward & landward margin of mangrove belt**.
- **'Opportunistic' colonization** due to ecological characteristics.
- Grows on **mud, sand, gravels, rocks, rock surfaces**.



# Adaptation = Development

- Modules for mangrove seedlings is moving to higher level of development for the villages, as **suitable materials have to be sourced, adapted & manufactured.**
- **Small machine shops** to produce modules in larger numbers & in standardized format, thus pushing development to a higher level. **More skills** to be developed as modules are then deployed to coasts.
- When mangroves are fully grown, forests themselves could open **opportunities for villages** to another level of development, e.g. **ecotourism development.**
- Timeline from mangrove seedlings to their maturity along coasts generates **increasing levels of development/opportunities** for villages & **at same time improves adaptation to SLR.**

# Winner of MIT Climate CoLab, 2014



WINNER

*Judges' Choice Award*


Poh Poh Wong

FOR THE PROPOSAL

**FUTURE MANGROVES**

IN THE 2014 ADAPTATION TO CLIMATE CHANGE CONTEST

**Professor Thomas W. Malone**  
Director, MIT Center for Collective Intelligence  
Patrick J. McGovern Professor of Management  
Sloan School of Management  
Massachusetts Institute of Technology

 **Climate CoLab**



# **Thank you**

**Email: [wong3921@gmail.com](mailto:wong3921@gmail.com)**

**Website: [coast4all](http://coast4all.com)**